

MEDICATIONS TO BE HELD FOR ALLERGY SKIN TESTING

Stop these medications **10** days prior to your appointment:

- Hydroxyzine(atarax®)
- *Doxepin
- *Imipramines
- Promethazine
- Meclizine
- Doxylamine
- Prochloroperazine

**Please discuss with your prescribing physician prior to stopping this medication.*

Stop these medications **7** days prior to your appointment:

- Allegra®, Allegra D® (Fexofenadine)
- Xyzal® (Levocetirizine)
- Zyrtec® (Cetirizine)
- Alavert® (Loratadine)
- Claritin®, Claritin D® (Loratadine)
- Clarinex® (Desloratadine)

Stop these medications **3** days prior to your appointment:

- Actifed® (Brompheniramine)
- Antihist®, Tavist® (Clemastine)
- Alka-Seltzer Plus Cold Medicine®
- Aller-Chlor®
- Ambenyl Cough Syrup®
- Astepro (Astelin®)
- Aventyl® (Nortriptyline)
- Benadryl® (Diphenhydramine)
- Bentlyl® (Dicyclomine)
- Bromfed®
- Chlortrimeton® (Chlorpheniramine)
- Citra Forte Syrup®
- Codimal®
- Comhist®
- Comtrex®
- Contac®
- Deconamine®
- Dimetapp® (Brompheniramine)
- Dramamine®
- Drixoral®
- Elavil® (Amitriptyline)
- Extendryl®
- Kronofed®
- Patanase®
- Pediacare®
- Percogesic®
- Periactin® (Cyproheptadine)
- Polaramine®
- Quelidrine®
- Rondec®
- Rynatan®
- Rynatuss®
- Scot-Tussin®
- Semprex D®
- Sinutab®
- Teldrin®
- Tigan®
- Triaminic®
- Tussionex®
- Tylenol PM®
- Xanax® (Alprazolam)
- 4-Way Cold Tablets®

Stop these medications **1** day prior to your appointment:

- Axid® (Nizatidine)
- Tagamet® (Cimetidine)

- Pepcid®
(Famotidine)
- Zantac® (Ranitidine)

Ocular (Eye) Drops:

- Ketotifen (Alaway®,
Zaditor®)
- Pataday®
- Patanol®
- Pazeo®
- Alcaftadine
(Lastacaft®)
- Azelastine (Optivar®)
- Bepostatine
(Bepreve®)
- Emadastine
(Emadine®)

Antihistamines are found in many medications (especially cold or allergy medications). If you do not know if your child's medicine contains antihistamines, ask the pharmacist or call the Allergy Clinic at **401-444-2489 or **401-444-6825**.**